

Umsichtig und vernünftig handeln auf der operativen Intensivstation

Autoren:

PD Dr. med. Rüdger Kopp

Klinik für Operative Intensivmedizin und Intermediate Care,
Uniklinik RWTH Aachen

Prof. Dr. med. Gernot Marx, FRCA

Klinik für Operative Intensivmedizin und Intermediate Care,
Uniklinik RWTH Aachen

Dr. med. René Wildenauer

Klinik und Poliklinik für Allgemein-, Viszeral, Gefäß- und Kinderchirurgie,
Universitätsklinikum Würzburg

Zitation:

Kopp R, Wildenauer R, Marx G: Umsichtig und vernünftig handeln auf der operativen Intensivstation. Passion Chirurgie. 2020 Oktober, 10(10): Artikel 04_02.

Literatur

- [1] Pronovost P, Berenholtz S, Dorman T, Lipsett PA, Simmonds T, Haraden C. Improving communication in the ICU using daily goals. *Journal of critical care*. 2003;18(2):71-5.
- [2] Lane D, Ferri M, Lemaire J, McLaughlin K, Stelfox HT. A systematic review of evidence-informed practices for patient care rounds in the ICU*. *Crit Care Med*. 2013;41(8):2015-29.
- [3] Berenholtz SM, Pronovost PJ, Lipsett PA, Hobson D, Earsing K, Farley JE, et al. Eliminating catheter-related bloodstream infections in the intensive care unit. *Crit Care Med*. 2004;32(10):2014-20.
- [4] Krishnan JA, Moore D, Robeson C, Rand CS, Fessler HE. A prospective, controlled trial of a protocol-based strategy to discontinue mechanical ventilation. *Am J Respir Crit Care Med*. 2004;169(6):673-8.
- [5] Kumpf O, Braun JP, Brinkmann A, Bause H, Bellgardt M, Bloos F, et al. Quality indicators in intensive care medicine for Germany - third edition 2017. *Ger Med Sci*. 2017;15:Doc10.
- [6] Reinhart K, Brunkhorst FM, Bone HG, Bardutzky J, Dempfle CE, Forst H, et al. Prävention, Diagnose, Therapie und Nachsorge der Sepsis. Erste Revision der S2k-Leitlinien der Deutschen Sepsis-Gesellschaft e.V. (DSG) und der Deutschen Interdisziplinären Vereinigung für Intensiv- und Notfallmedizin (DIVI). *Der Anaesthesist*. 2010;59(4):347-70.
- [7] Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Med*. 2017;43(3):304-77.
- [8] Azuhata T, Kinoshita K, Kawano D, Komatsu T, Sakurai A, Chiba Y, et al. Time from admission to initiation of surgery for source control is a critical determinant of survival in patients with gastrointestinal perforation with associated septic shock. *Crit Care*. 2014;18(3):R87.

- [9] Bloos F, Thomas-Ruddel D, Ruddel H, Engel C, Schwarzkopf D, Marshall JC, et al. Impact of compliance with infection management guidelines on outcome in patients with severe sepsis: a prospective observational multi-center study. *Crit Care*. 2014;18(2):R42.
- [10] Solomkin JS, Mazuski JE, Bradley JS, Rodvold KA, Goldstein EJ, Baron EJ, et al. Diagnosis and management of complicated intra-abdominal infection in adults and children: guidelines by the Surgical Infection Society and the Infectious Diseases Society of America. *Clin Infect Dis*. 2010;50(2):133-64.
- [11] Marx G, Schindler AW, Mosch C, Albers J, Bauer M, Gnass I, et al. Intravascular volume therapy in adults: Guidelines from the Association of the Scientific Medical Societies in Germany. *Eur J Anaesthesiol*. 2016;33(7):488-521.
- [12] Monnet X, Marik P, Teboul JL. Passive leg raising for predicting fluid responsiveness: a systematic review and meta-analysis. *Intensive Care Med*. 2016;42(12):1935-47.
- [13] Marik PE, Cavallazzi R. Does the central venous pressure predict fluid responsiveness? An updated meta-analysis and a plea for some common sense. *Crit Care Med*. 2013;41(7):1774-81.
- [14] Guay J, Ochroch EA, Kopp S. Intraoperative use of low volume ventilation to decrease postoperative mortality, mechanical ventilation, lengths of stay and lung injury in patients without acute lung injury. *Cochrane Database Syst Rev*. 2018;7:CD011151.
- [15] Neto AS, Simonis FD, Barbas CS, Biehl M, Determann RM, Elmer J, et al. Lung-Protective Ventilation With Low Tidal Volumes and the Occurrence of Pulmonary Complications in Patients Without Acute Respiratory Distress Syndrome: A Systematic Review and Individual Patient Data Analysis. *Crit Care Med*. 2015;43(10):2155-63.
- [16] Fan E, Del Sorbo L, Goligher EC, Hodgson CL, Munshi L, Walkey AJ, et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. *Am J Respir Crit Care Med*. 2017;195(9):1253-63.
- [17] Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA*. 2012;307(23):2526-33.
- [18] Bein T, Bischoff M, Bruckner U, Gebhardt K, Henzler D, Hermes C, et al. S2e guideline: positioning and early mobilisation in prophylaxis or therapy of pulmonary disorders : Revision 2015: S2e guideline of the German Society of Anaesthesiology and Intensive Care Medicine (DGAI). *Der Anaesthesist*. 2015;64 Suppl 1:1-26.
- [19] S3-Leitlinie Invasive Beatmung und Einsatz extrakorporaler Verfahren bei akuter respiratorischer Insuffizienz: Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften, AWMF; 2017 [Available from: <https://www.awmf.org/leitlinien/detail/II/001-021.html>], abgerufen am 08.11.2017.
- [20] Schaller SJ, Anstey M, Blobner M, Edrich T, Grabitz SD, Gradwohl-Matis I, et al. Early, goal-directed mobilisation in the surgical intensive care unit: a randomised controlled trial. *Lancet*. 2016;388(10052):1377-88.
- [21] Bingold T, Bickenbach J, Coburn M, David M, Dembinski R, Kuhnle G, et al. Modulares Zertifikat Intensivmedizin der DGAI. *Anasth Intensivmed*. 2014;55:316-29.
- [22] Wacha H, Hoyme U, Isenmann R, Kujatz P, Lebert C, Naber K, et al. Perioperative Antibiotika-Prophylaxe. Empfehlungen einer Expertenkommission der Paul-Ehrlich-Gesellschaft für Chemotherapie e. V. *Chemother J*. 2010;19(3):70-84.
- [23] Reutter F, Reuter DA, Hilgarth H, Heilek AM, Goepfert MS, Punke MA. Perioperative Antibiotikaprophylaxe. *Der Anaesthesist*. 2014;63(1):73-86.
- [24] WHO Guidelines Approved by the Guidelines Review Committee. *Global Guidelines for the Prevention of Surgical Site Infection*. Geneva: World Health Organization. Copyright (c) World Health Organization 2016.; 2016.
- [25] Yunos NM, Bellomo R, Glassford N, Sutcliffe H, Lam Q, Bailey M. Chloride-liberal vs. chloride-restrictive intravenous fluid administration and acute kidney injury: an extended analysis. *Intensive Care Med*. 2015;41(2):257-64.

- [26] Yunos NM, Bellomo R, Hegarty C, Story D, Ho L, Bailey M. Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA*. 2012;308(15):1566-72.
- [27] Semler MW, Self WH, Wanderer JP, Ehrenfeld JM, Wang L, Byrne DW, et al. Balanced Crystalloids versus Saline in Critically Ill Adults. *N Engl J Med*. 2018;378(9):829-39.
- [28] Chant C, Wilson G, Friedrich JO. Anemia, transfusion, and phlebotomy practices in critically ill patients with prolonged ICU length of stay: a cohort study. *Crit Care*. 2006;10(5):R140.
- [29] Kumwilaisak K, Noto A, Schmidt UH, Beck CI, Crimi C, Lewandrowski K, et al. Effect of laboratory testing guidelines on the utilization of tests and order entries in a surgical intensive care unit. *Crit Care Med*. 2008;36(11):2993-9.
- [30] Ganapathy A, Adhikari NK, Spiegelman J, Scales DC. Routine chest x-rays in intensive care units: a systematic review and meta-analysis. *Crit Care*. 2012;16(2):R68.
- [31] Muller MCA, Stanworth SJ, Coppens M, Juffermans NP. Recognition and Management of Hemostatic Disorders in Critically ill Patients Needing to Undergo an Invasive Procedure. *Transfusion medicine reviews*. 2017;31(4):223-9.
- [32] Warner MA, Chandran A, Jenkins G, Kor DJ. Prophylactic Plasma Transfusion Is Not Associated With Decreased Red Blood Cell Requirements in Critically Ill Patients. *Anesth Analg*. 2017;124(5):1636-43.
- [33] Hebert PC, Carson JL. Transfusion threshold of 7 g per deciliter--the new normal. *N Engl J Med*. 2014;371(15):1459-61.
- [34] Bundesärztekammer. Querschnitts-Leitlinien zur Therapie mit Blutkomponenten und Plasmaderivaten. 4 ed: Deutscher Ärzte-Verlag; 2015.
- [35] Bienholz A, Kribben A. KDIGO-Leitlinien zum akuten Nierenversagen. *Der Nephrologe*. 2013;8(3):247-51.
- [36] Ho KM, Sheridan DJ. Meta-analysis of frusemide to prevent or treat acute renal failure. *BMJ (Clinical research ed)*. 2006;333(7565):420.
- [37] Fliser D, Laville M, Covic A, Fouque D, Vanholder R, Juillard L, et al. A European Renal Best Practice (ERBP) position statement on the Kidney Disease Improving Global Outcomes (KDIGO) clinical practice guidelines on acute kidney injury: part 1: definitions, conservative management and contrast-induced nephropathy. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2012;27(12):4263-72.
- [38] Händehygiene in Einrichtungen des Gesundheitswesens Empfehlung der Kommission für Krankenhaushygiene und Infektionsprävention (KRINKO) beim Robert Koch-Institut (RKI). *Bundesgesundheitsbl*. 2016;59:1189-220.
- [39] Scheithauer S, Haefner H, Schwanz T, Schulze-Steinen H, Schiefer J, Koch A, et al. Compliance with hand hygiene on surgical, medical, and neurologic intensive care units: direct observation versus calculated disinfectant usage. *Am J Infect Control*. 2009;37(10):835-41.
- [40] SepNet Critical Care Trials G. Incidence of severe sepsis and septic shock in German intensive care units: the prospective, multicentre INSEP study. *Intensive Care Med*. 2016;42(12):1980-9.