

## „The big Four“ – BCC, PEC, MM & MCC: Stadiengerechte Therapie von Hautmalignomen der Kopf-Hals-Region.

### Korrespondierender Autor:

PD Dr. med. Dr. med. dent. Jan D. Raguse

Fachklinik Hornheide

Klinik für Mund-, Kiefer- und Gesichtschirurgie/ Plastische Operationen

Dorbaumstrasse 300

48157 Münster

raguse@fachklinik-hornheide.de

### Zitation:

Raguse J-D, K. Wermker K: „The big Four“ – BCC, PEC, MM & MCC: Stadiengerechte Therapie von Hautmalignomen der Kopf-Hals-Region. Passion Chirurgie. 2021 April; 11(04): Artikel 03\_03.

### Literatur

- [1] Lang BM, Balermipas P, Bauer A, Blum A, Brölsch GF, Dirschka T, et al. S2k Guidelines for Cutaneous Basal Cell Carcinoma - Part 2: Treatment, Prevention and Follow-up. J Dtsch Dermatol Ges J Ger Soc Dermatol JDDG. 2019;17:214–30.
- [2] Lang BM, Grabbe S. [Diagnosis and treatment of basal cell carcinoma : A question of the risk of recurrence]. Hautarzt Z Dermatol Venerol Verwandte Geb. 2020;71:580–7.
- [3] Eberle FC, Kanyildiz M, Schnabl SM, Schulz C, Häfner H-M, Adam P, et al. Three dimensional (3D) histology in daily routine: practical implementation and its evaluation. J Dtsch Dermatol Ges J Ger Soc Dermatol JDDG. 2014;12:1028–35.
- [4] Leiter U, Heppt MV, Steeb T, Amaral T, Bauer A, Becker JC, et al. S3 guideline for actinic keratosis and cutaneous squamous cell carcinoma (cSCC) - short version, part 2: epidemiology, surgical and systemic treatment of cSCC, follow-up, prevention and occupational disease. J Dtsch Dermatol Ges J Ger Soc Dermatol JDDG. 2020;18:400–13.
- [5] Heppt MV, Leiter U, Steeb T, Amaral T, Bauer A, Becker JC, et al. S3 guideline for actinic keratosis and cutaneous squamous cell carcinoma - short version, part 1: diagnosis, interventions for actinic keratoses, care structures and quality-of-care indicators. J Dtsch Dermatol Ges J Ger Soc Dermatol JDDG. 2020;18:275–94.
- [6] Gutzmer R, Wiegand S, Kölbl O, Wermker K, Heppt M, Berking C. Actinic Keratosis and Cutaneous Squamous Cell Carcinoma. Dtsch Arzteblatt Int. 2019;116:616–26.
- [7] Wermker K. Metastasierungsweg und Lymphknotenausräumung beim Unterlippenkarzinom. MKG-Chir. 2018;11:103–10.
- [8] Wermker K, Kluwig J, Schipmann S, Klein M, Schulze HJ, Hallermann C. Prediction score for lymph node metastasis from cutaneous squamous cell carcinoma of the external ear. Eur J Surg Oncol J Eur Soc Surg Oncol Br Assoc Surg Oncol. 2015;41:128–35.

- [9] Wermker K, Belok F, Schipmann S, Klein M, Schulze H-J, Hallermann C. Prediction model for lymph node metastasis and recommendations for elective neck dissection in lip cancer. *J Cranio-Maxillo-fac Surg Off Publ Eur Assoc Cranio-Maxillo-fac Surg.* 2015;43:545–52.
- [10] National Cancer Registration and Analysis Service. Rare skin cancer in England. Available at: [http://www.ncin.org.uk/publications/data\\_briefings/rareskincancer](http://www.ncin.org.uk/publications/data_briefings/rareskincancer) (accessed 26 October 2020)
- [11] Becker JC, Stang A, Decaprio JA, et al. Merkel cell carcinoma. *Nat Rev Dis Primers* 2017;3:17077. Heath M, Jaimes N, Lemos B et al. Clinical characteristics of Merkel cell carcinoma at diagnosis in 195 patients: the AEIOU features. *J Am Acad Dermatol* 2008; 58: 375–81.
- [12] Feng H, Shuda M, Chang Y, Moore PS. Clonal integration of a polyomavirus in human Merkel cell carcinoma. *Science* 2008;319:1096–1100.
- [13] Agelli M, Clegg LX. Epidemiology of primary Merkel cell carcinoma in the United States. *J Am Acad Dermatol* 2003; 49: 832–41.
- [14] Coggshall K, Tello TL, North JP, Yu SS. Merkel cell carcinoma: an update and review: pathogenesis, diagnosis, and staging. *J Am Acad Dermatol* 2018; 78: 433–42.
- [15] Vandeven N, Lewis CW, Makarov V, et al. Merkel cell carcinoma patients presenting without a primary lesion have elevated markers of immunity, higher tumor mutation burden, and improved survival. *Clin Cancer Res* 2018;24:963–71.
- [16] Fondain M, Dereure O, Uhry Z, et al. Merkel cell carcinoma in France: a registries-based, comprehensive epidemiological survey. *J Eur Acad Dermatol Venereol* 2018 Jan 16. doi: 10.1111/jdv.14798. [Epub ahead of print]
- [17] Sridharan V, Muralidhar V, Margalit DN, et al. Merkel cell carcinoma: A population analysis on survival. *J Natl Compr Canc Netw* 2016;14:1247–57.
- [18] Frohm ML, Griffith KA, Harms KL, et al. Recurrence and survival in patients with Merkel cell carcinoma undergoing surgery without adjuvant radiation therapy to the primary site. *JAMA Dermatol* 2016;152:1001–7.
- [19] Fields RC, Busam KJ, Chou JF, et al. Recurrence after complete resection and selective use of adjuvant therapy for stage I through III Merkel cell carcinoma. *Cancer* 2012;118:3311–3320.
- [20] Kang SH, Haydu LE, Goh RY, Fogarty GB. Radiotherapy is associated with significant improvement in local and regional control in Merkel cell carcinoma. *Radiat Oncol* 2012;7:171.
- [21] S2k-Guideline Merkel Cell Carcinoma (Registernr.: 032-023) - Update 2018
- [22] Hoeller U, Mueller T, Schubert T, et al. Regional nodal relapse in surgically staged Merkel cell carcinoma. *Strahlenther Onkol.* 2015;191:51–8.
- [23] Kouzmina M, Leikola J, Böhling T, Koljonen V. Positive sentinel lymph node biopsy predicts local metastases during the course of disease in Merkel cell carcinoma. *J Plast Surg Hand Surg* 2013;47:139–43.
- [24] Nghiem P, Kaufman HL, Bharmal M, et al. Systematic literature review of efficacy, safety and tolerability outcomes of chemotherapy regimens in patients with metastatic Merkel cell carcinoma. *Future Oncol* 2017;13:1263–79.
- [25] Iyer JG, Blom A, Doumani R, et al. Response rates and durability of chemotherapy among 62 patients with metastatic Merkel cell carcinoma. *Cancer Med* 2016;5:2294–301.
- [26] Kaatsch P., et al., *Krebs in Deutschland 2011/2012.* 2015, Robert-Koch-Institut: Berlin. <http://www.rki.de>
- [27] Saarland, K. 2012 [cited 2012; Available from: <http://www.krebsregister.saarland.de/>.

- [28] <https://www.krebsgesellschaft.de/onko-internetportal/basis-informationen-krebs/krebsarten/hautkrebs/malignes-melanom-schwarzer-hautkrebs.html>
- [29] Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF): S3-Leitlinie Prävention von Hautkrebs, Langversion 1. 1, 2014, AWMF Registernummer: 032/052OL, <http://leitlinienprogramm-onkologie.de/Leitlinien.7.0.html>
- [30] Arbeitsgemeinschaft Dermatologische Prävention; <https://www.unserehaut.de>
- [31] Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, 12 AWMF): Diagnostik, Therapie und Nachsorge des Melanoms, Langversion 3.2, 2019, AWMF Registernummer: 032/024OL, <http://www.leitlinienprogramm-onkologie.de/leitlinien/melanom/>
- [32] Tran, K.T., N.A. Wright, and C.J. Cockerell, Biopsy of the pigmented lesion--when and how. *J Am Acad Dermatol*, 2008. 59(5): p. 852-71
- [33] Wheatley, K., et al., Surgical excision margins in primary cutaneous melanoma: A meta-analysis and Bayesian probability evaluation. *Cancer Treat Rev*, 2016. 42: p. 73-81.
- [34] McLeod, M., et al., Surgical treatments for lentigo maligna: a review. *Dermatol Surg*, 2011. 37(9): p. 1210-28. <https://www.ncbi.nlm.nih.gov/pubmed/21631635>
- [35] Mohrle, M., et al., [Prognostic parameters and surgical strategies for facial melanomas]. *J Dtsch Dermatol Ges*, 2003. 1(6): p. 457-63.
- [36] van der Ploeg, A.P., et al., Prognosis in patients with sentinel node-positive melanoma is accurately defined by the combined Rotterdam tumor load and Dewar topography criteria. *J Clin Oncol*, 2011. 29(16): p. 2206-14.
- [37] Frankel, T.L., et al., Do micromorphometric features of metastatic deposits within sentinel nodes predict nonsentinel lymph node involvement in melanoma? *Ann Surg Oncol*, 2008. 15(9): p. 2403-11
- [38] Medina, J.E., et al., Current management of cutaneous malignant melanoma of the head and neck. *Acta Otolaryngol*, 2002. 122(8): p. 900-6.
- [39] Schmalbach, C.E., T.M. Johnson, and C.R. Bradford, The management of head and neck melanoma. *Curr Probl Surg*, 2006. 43(11): p. 781-835.
- [40] Wiener, M., R.F. Uren, and J.F. Thompson, Lymphatic drainage patterns from primary cutaneous tumours of the forehead: refining the recommendations for selective neck dissection. *J Plast Reconstr Aesthet Surg*, 2014. 67(8): p. 1038-44.
- [41] Gründahl JEH, Hallermann C, Schulze H-J, Klein M, Wermker K. Cutaneous Angiosarcoma of Head and Neck: A New Predictive Score for Locoregional Metastasis. *Transl Oncol*. 2015;8:169–75.